1x RBC Lysis Buffer (250ml)

Order No.:60-00050-12

Description

Red Blood Cell (RBC) Lysis Buffer has been designed, formulated, and tested to ensure optimal lysis of RBCs in single cell suspensions with minimal effects on leukocytes. Nucleated RBCs are not effectively lysed with ammonium chloride. This 1x RBC Lysis Buffer is ready to use.

Components

- NH 4 Cl (155 mMol/L)
- KHCO 3 (10 mMol/L)
- EDTA (0,1 mMol/L)

Protocol/h3>

1. **Ensure that the RBC Lysis Buffer is cold (4-8°C) .**
2. **Dilute one volume of cell suspension with 10 volumes of 1× Red Blood Cell Lysis Solution.**
3. ***When working wih blood samples:***
   Vortex for 5 seconds and incubate for 10-15 minutes (no more than 15 minutes) at 4-8°C.

   When working with tissue samples:
   Vortex for 5 seconds and incubate for 2 minutes at 4-8°C.
   Note: Observe turbidity to evaluate red blood cell lysis. Once the sample becomes clear red, lysis is complete.
4. **Vortex for 5 seconds and centrifuge at 300×g for 10 minutes at room temperature.**
5. **Aspirate supernatant completely.**
6. **Resuspend the cell pellet in an appropriate buffer and proceed to further applications.**

pluriSelect Life Science
Deutscher Platz 5a, 04103 Leipzig, DE
Phone: +49 341 333 858 0 | Fax: +49 341 333 858 99
support@pluriselect.com
https://www.pluriselect.com/

Additional Information

Size	250 ml
Delivery Time (days)	1-2
Area of interest	Cell Biology
Sterility	Sterile
pH Range	pH 7.4
Shipping Condition	Room Temperature
Storage Condition	4-8 °C
Regulatory Statement	For research use only. Not for use in diagnostic procedures.,

Warning and Limitations

This product is for research and development only, not for diagnostic or theurapeutic use.

pluriSelect Life Science
Deutscher Platz 5a, 04103 Leipzig, DE
Phone: +49 341 333 858 0 | Fax: +49 341 333 858 99
support@pluriselect.com
https://www.pluriselect.com/