

## 1x RBC Lysis Buffer (100ml)

Order No.:60-00050-11



### Description

Red Blood Cell (RBC) Lysis Buffer has been designed, formulated, and tested to ensure optimal lysis of RBCs in single cell suspensions with minimal effects on leukocytes. Nucleated RBCs are not effectively lysed with ammonium chloride. This 1x RBC Lysis Buffer is ready to use.

### Components

- NH<sub>4</sub>Cl (155 mMol/L)
- KHCO<sub>3</sub> (10 mMol/L)
- EDTA (0,1 mMol/L)

### Protocol

1. Ensure that the RBC Lysis Buffer is cold (4-8°C).
2. Dilute one volume of cell suspension with 10 volumes of 1x Red Blood Cell Lysis Solution.
3. *When working with blood samples:*  
Vortex for 5 seconds and incubate for 10-15 minutes (no more than 15 minutes) at 4-8°C.

*When working with tissue samples:*

Vortex for 5 seconds and incubate for 2 minutes at 4-8°C.

**Note:** Observe turbidity to evaluate red blood cell lysis. Once the sample becomes clear red, lysis is complete.

4. Vortex for 5 seconds and centrifuge at 300×g for 10 minutes at room temperature.
5. Aspirate supernatant completely.
6. Resuspend the cell pellet in an appropriate buffer and proceed to further applications.

### Additional Information

Size	100 ml
Delivery Time (days)	1-2

#### PluriSelect USA

2030 Gillespie Way, Unit 103, CA 92020 El Cajon, US  
Phone: +1 619-202-4297 | Fax: +1 619-828-8003  
sales-support@pluriselect-usa.com  
<https://www.pluriselect.com/>

Area of interest	Cell Biology
Sterility	Sterile
pH Range	pH 7.4
Shipping Condition	Room Temperature
Storage Condition	4-8 °C
Regulatory Statement	For research use only. Not for use in diagnostic procedures.,

### **Warning and Limitations**

This product is for research and development only, not for diagnostic or therapeutic use.