

pluriStrainer® 50 μm (Cell Strainer)



Order No.	Sterlity	Packaging	Variant
43-50050-03	Sterile	Individual single packed	25 pcs, sterile
43-50050-50	Non-Sterile	50 pcs. / Bag	50 pcs, non-sterile

Description

The pluriStrainer® is a sieving device (or cell strainer) for a wide range of laboratory applications where the filtration and the purification of liquids is required. Its unique design features improved ventilation to avoid clogging for a smart sample preparation. pluriStrainer® are stackable to allow direct filtration with different mesh sizes and alsocan be inverted to recover the sieved material. For large sample volumes it can be combined with a funnel. It fits all major brand 50 mL centrifuge (conical) tubes. The combination of the pluriStrainer® and the Connector Ring allow using low pressure which supports the filtration. Furthermore it is possible to block the flow through the sieve and hold back the liquid on top of the mesh. The liquid on top of the mesh can be used for sample preparation and incubation such as tissue dissociation or the treatment of cells with cell lysis reagents (e.g. Buffer RLT, TRIzol®).

Application

- Obtaining a real single cell suspension after digestion of mammary tissue and organoids
- Enrichment of specific cells using pluriBead technology
- Preparation of a single cell suspension of blood cells from bone marcol-sm-12, pancreas, thymus, lymph nodes and others
- Dissociation of cells from other primary tissue
- Preparation of real single cell suspension for flow cytometry(FACS™)
- Faster and easier alternative to gauze filtration
- Sieving of complex and viscous liquids (in combination with Connector Ring & Syringe for vacuum)
- Short time incubation of cells / cytokine (in combination with pluriBead and Connector Ring)

Features



Filter
Attach pluriStrainer® to a sterile 50 mL centrifuge tube. Then, add sample material onto the strainer and filter sample



Reverse

To obtain the larger fraction, take off the pluriStrainer®, turn it upside down onto another 50 mL tube and flush back the sample from the pluriStrainer®.



Increase sample load

With Funnel You can add up to 24 mL sample material on top.



Stack $Stacking \ of \ pluriStrainer @\ with \ different \ mesh \ sizes \ allows \ for \ straining \ various \ cell \ sizes \ at \ the \ same \ time.$



Flow control

Allows to control the rate of flow by opening or closing the Luer-Lock, e.g. for physical dissociation of primary tissue (brain, spleen etc.).



Low pressure

If you add a syringe to the Connector Ring, it is possible to force low pressure to support the straining of rough sample material while pulling the piston.

Additional Information

Delivery Time (days)	1-2
Mesh Size	50 μm
Packaging	Individual single packed
Color	gray
Flow Control	Yes, in combination with Connecting Ring
Mesh Fixation	Injected in housing for strong hold
Fabric material	PET (Polyethylenterephthalat)

Housing Material	LD-PE (Low Density Polyethylen)	
Mesh Type	woven	
Handling	Extended lip on strainer enables aseptic handling	
Sterility	Sterile	
Comparable with	Corning Cell Strainer, EASYstrainer™ Cell sieve, Falcon™ cell strainer, Reversible Strainer, MACS® SmartStrainer	
Stability	Buffer RLT®, TRIzol®, Isopropanol, organic solvent	
Centrifugable	Yes	
Tissue Dissociation	Yes	
Disinfectable	Yes, with 70% ethanol	
Area of interest	Immunology, Cell Biology, Plant Biology, Microbiology, Oncology, Ecology	
Shipping Condition	Room Termperature	
Storage Condition	Room Temperature	
Regulatory Statement	For research use only. Not for use in diagnostic procedures,	
Legal information	Buffer RLT® is a trademark of Qiagen, TRIzol® is a trademark of Molecular Research Center, Inc., FACS™ is a trademark of BD Biosciences, Corning is a registered trademark of Corning, Inc.; EASYstrainer™ is a registered trademark of Greiner Bio One International GmbH, Falcon™ Cell Strainers is a registered trademark of Corning, Inc.; MACS® SmartStrainer is a registered trademark of Miltenyi Biotec GmbH	

Warning and Limitations

This product is for research and development only, not for diagnostic or theurapeutic use.